



Your source for the latest research news

One Drop. Once a Day.
New PATADAY™ solution
The most common side effects include cold, sore throat, blurred vision, burning or stinging.

Visit our site for a \$10 rebate on New PATADAY™ solution.

NEW! Once-a-Day Pataday™
(olopatadine hydrochloride ophthalmic solution) 0.2%



Latest News

Browse Topics

Encyclopedia

Health Center

Videos

Science Books

Careers

Health & Medicine

Mind & Brain

Plants & Animals

Space & Time

Earth & Climate

Matter & Energy

Computers & Math

Fossils & Ruins

Show menu

ScienceDaily home page

RSS feeds

Free newsletter

Print this page

Email to friend

Bookmark

Text: small | med | large

Source: [Universidad de Granada](#)
Date: May 9, 2007

More on: [Infant's Health](#), [Pregnancy and Childbirth](#), [Disability](#), [Birth Defects](#), [Gynecology](#), [Down's Syndrome](#)

Wait A Few Minutes Before Clamping The Umbilical Cord, Researchers Suggest

Science Daily — In the past, the newborns' umbilical cord was not clamped right after birth, thus allowing the blood flow to stop naturally. This practice, known as “late clamping”, was replaced by “early clamping”, that is, cutting the cord immediately after the infant is expelled. However, this new practice lacks studies corroborating its benefits. In fact, recent studies on the importance of when clamping should be done have shown contradictory results.

A doctoral thesis carried out at the Department of Obstetrics and Gynecology at the University of Granada by Catalina de Paco Matallana shows that the clamping of the umbilical cord of newborns from full-term pregnancies (that is, infants born after a nine-month pregnancy) two minutes after the infant is expelled from the womb makes no difference to hematocrit or hemoglobin levels of the umbilical cord vein compared to clamping the cord within 20 seconds. Thus, the study shows that early clamping (which is widely performed) is not justified.

Ads by Google

Advertise on this site

Baby stem cell storage
The experts in cord blood stem cell collection & storage
www.cells4life.co.uk

Cerebral Palsy Treatment
New research and treatments being done with cord blood stem cells
www.stemcellschina.com

Baby's Ultrasound Images
Your free weekly pregnancy guide, fetal ultrasound images, and more.
pregnancyweekly.com

Costa del Sol Midwife
Pregnant? Need help and support? You need an English Midwife!
www.costadelsolmidwife.com

Pregnancy Information
Answers to your questions about pregnancy, labor and more!
obgyn.health.ivillage.com

Analysis of 151 umbilical cords

The doctoral thesis *Repercusiones clínicas y fisicoquímicas del tiempo de ligadura del cordón umbilical en recién nacidos a término* (Clinical and Physiochemical Repercussions of Clamping Time of the Umbilical Cord of Newborns from Full-Term Pregnancies) analyzed a total of 151 umbilical cords of newborns from full-term pregnancies. In 79 cases, the umbilical cord was cut within 20 seconds, and in 72 cases it was clamped two minutes after the infant was expelled.

The study found that the partial pressure of oxygen in the umbilical artery of the newborn babies who had late clamping had risen, while there was a lesser need of oxygenotherapy after birth. There were no differences in the removal time of the placenta and the mother's bleeding after birth - one of the reasons why early clamping of the umbilical cord started to be practiced.


De Paco Matallana also analyzed the melatonin concentration (one of the strongest and most currently studied antioxidants), finding differences between the melatonin concentration (aMT) in the umbilical vein and the umbilical artery, the latter being where the concentration is significantly higher. “This suggests that the fetus not only receives melatonin from the mother via the umbilical vein because it crosses the placental barrier, but also that it is produced in stressful situations such as during labor,” says de Paco Matallana. Thus, high melatonin concentrations in the umbilical cord, together with the arterial and vein differences according to type of birth, suggest that the pineal gland is used by the fetus and that it can respond during birth.

Preventing anemia in children

As for the concentration of the triacylglycerols (TAG) analyzed in the umbilical vein of both groups, this study found statistically significant differences, as there is a higher concentration of TAG in the group of early clamping. The same is applied to the umbilical artery, where there are also statistically significant differences, with higher a concentration of TAG in the group of early clamping. However, there are no studies corroborating these findings, so more research is needed, although many other studies recommend late clamping “especially because of the beneficial effect on the prevention of anemia in children,” says de Paco Matallana.

The author points out that the results of her thesis “show that there are no differences in the hemogram or the general biochemical profile in the umbilical vein in the cases of early and late clamping”. Moreover, there were no

Health Videos & Features



Helping Kids With ADHD Succeed in School
Learn how to help children with attention deficit hyperactivity disorder (ADHD) improve their school performance, with medical and behavior therapies.

- News:
- Treating Teens Who Feel the Burn of GERD
 - Keeping Chickenpox All Cooped Up
 - Injury Prevention Outside the Home for One- to Two-Year-Olds

Baby stem cell storage
The experts in cord blood stem cell collection & storage
www.cells4life.co.uk

Granada Airport car hire
Ford Ka from £53 week Car Hire of the year 2006
www.carjet.co.uk

Ads by Google

Fertility FACT 27

Infertility is due to male factors in about 35% of cases.

The good news? Treatments are available for many conditions.

Be informed. Take action. Visit fertilityjourney.com

Related News Sections

- [Health & Medicine](#)

Related News Topics

- [Infant's Health](#)
 - [Pregnancy and Childbirth](#)
 - [Disability](#)
- [Birth Defects](#)
 - [Gynecology](#)
 - [Down's Syndrome](#)

Related Science Stories

- [Delayed Umbilical Cord Clamping Boosts Iron In Infants](#)
- [Topical Antiseptic Reduces Umbilical Cord Infection And Mortality Risk](#)
- [Newborn Brain Injuries Stem From Infections, Not Delivery](#)
- [New Stem Cell Source Could Boost Bone Marrow Success; Uncharted Area Of Umbilical Cord Offers Hope](#)
- [Leukemia Patients Survive With Stem Cell Transplant](#)

Related Encyclopedia Articles

- [Umbilical cord](#)
 - [Placenta](#)
 - [Infant](#)
 - [Childbirth](#)
 - [Fetus](#)
- [Stillbirth](#)
 - [Breech birth](#)
 - [Stem cell](#)
 - [Birth weight](#)
 - [Vocal folds](#)

Related Book Reviews

- [Patch Clamping : An Introductory Guide to Patch Clamp Electrophysiology](#)
- [Weight Watchers Make It in Minutes: Easy](#)

Search Archives

Find:

Go > options

JobStream

Quick Search


Material Science Engineer
Think Energy Group
US-TX-Austin


Manager, Food Science, Food, beverage, product development
CyberCoders Engineering
US-NY-Ithaca

Director Bioanalytical Science - Chemistry - LC/MS Methods
CyberCoders - Biotechnology, Life Science and Pharmaceutical

careerbuilder.com

Now in paperback





In Other News ...

- Giuliani to affirm abortion-rights support** (6 hours ago)

Sparks expected to fly in Padilla trial (6 hours ago)

Troops moved from Manila before elections (6 hours ago)

Norwegian meat packer recalls ham (6 hours ago)

Girl may go to Britain for abortion

differences in the viscosity or the melatonin in the umbilical artery and vein, which have traditionally been variables related to early clamping. From a clinical point of view, there were no differences in either group with umbilical clamping at birth.

“This study has not found any scientific evidence to suggest that the practice of early clamping is advisable or to justify the abandonment of late clamping in newborns from full-term pregnancies,” says de Paco Matallana. “There are convincing findings for and against the two different types of clamping analyzed in this study, which shows not only the complexity of the problem, but also that research in this field may not be controlled enough or designed correctly.”

Reference: Catalina de Paco Matallana. Department of Obstetrics and Gynecology of the University of Granada.

Note: This story has been adapted from a news release issued by Universidad de Granada.

Ads by Google

Advertise on this site

Treat Language Delays Unlock your child's potential for speech and language with our DVDs www.babybumblebee.com	Childbirth 2006 Webby Award Winner-Best Health Website. Doctor Articles & More! www.healthline.com	Moms Skin Rash Cure Mother discovers product to heal any skin rash in 48 hours. Amazing www.freederm.com
---	---	---

New! Search [Science Daily](#) or the entire web with Google:



☐ Web ☒ ScienceDaily.com

- [Recipes in 15, 20, and 30 Minutes](#)
- [Childhood Brain & Spinal Cord Tumors: A Guide for Families, Friends & Caregivers](#)
 - [Ajax in Action](#)
 - [The South Beach Diet Quick and Easy Cookbook : 200 Delicious Recipes Ready in 30 Minutes or Less](#)

(6 hours ago)

III. lawmakers seek ban on horse slaughter (6 hours ago)

Man kills mother, self with saw (7 hours ago)

Britain forces all-white schools to mix (7 hours ago)

Cracker Barrel brings burgers back (7 hours ago)

Federal education mandates faulted (7 hours ago)

... more breaking news at **NewsDaily** -- updated every 15 minutes