Forgot Password New Us



About Us | Contact Us | Feedback | Portfolio | Press | Advertise | Careers | Sitemap | Log In

Baby stem cell collection

Granada Airport car hire The experts in stem cell cord blood collection Ford Ka from £53 week Car Hire of the year 2006

Login Password

Forums

Umbilical Delivery

Google SEARCH

Home **Directories Professionals Consumers Education Business** News **Websites** Web Medindia Search Advanced Search XML Health news XML Disease-News

Ads by Google **Baby Umbilical Cord** Cathedral of Granada **Umbilical Pregnancy Umbilical Blood**

Medindia » Latest Health News » Late Clamping of Umbilical Cord may Reduce Anemia in Children

Latest Headlines

Statins may Help Treat Hepatitis C Patients

Sleep Apnea Patients Have Greatly Increased Risk of Severe Car Crashes

Now, Cellphones That Alert Against Impending Lightning Strikes

Start School Later in the Morning, Say Sleepy Teens

'wake Up and Breathe' Strategy Allows Patients to Come Off Ventilator Sooner

Veterans Exposed to Agent Orange Increased Risk of Prostate Cancer Recurrence

Green Tea may Protect the Bladder from Becoming Inflamed

Genetic Marker Linked to Aggressive Prostate Cancer

Drug Eluting Nanostructured Coatings Enable Targeted Drug Therapy for Orthopedic Patients

Nanomedicine Gives the Way for Nerve Cell Regeneration

Health Guide A-Z

Health Topics A-Z

News A-Z

Diseases A-Z

First Aid A-Z

Drug A-Z

Procedures A-Z

Diet & Nutrition A-Z

Drug Toxicity A-Z

Lab Tests A-Z

Animation A-Z

Calculators A-Z

Glossary A-Z

All Latest News

Child Health News

Health News Posted on May 17, 2007

🔀 Email 🔝 RSS Feeds 🛭 🖨 Print 🔚 Save 🦺 Link 🥞 Syndicate 📝 Comments

Late Clamping of Umbilical Cord may Reduce Anemia in Children

In the past, the newborns' umbilical cord was not clamped right after birth, thus allowing the blood flow to stop naturally. This practice, known as "late clamping", was replaced by "early clamping", that is,

cutting the cord immediately after the infant is expelled. However, this new practice lacks studies corroborating its benefits. In fact, recent studies on the importance of when clamping should be done have shown contradictory results.

Cord Blood Stem Cells

Granada

A doctoral thesis carried Store your child's cord blood stem cells with Cryo-Save www.cryo-save.com out at the Department of Obstetrics and Gynecology

of Visiting Granada? Find Deals & Read Hotel Reviews! University Granada by Catalina Paco Matallana shows that 50 Hotels in Granada

the map! clamping of umbilical cord of newborns from full-term pregnancies Property in Granada

(that is, infants born after a nine-month pregnancy) two minutes after the infant is

www.bookings.es

www.TripAdvisor.com

Andalusia's last secret. Culture of white-washed villages. www.qualitv-sol.com

Book your hotel in Granada online. Find your hotel on a city

Ads by Google

expelled from the womb makes no difference to hematocrit or hemoglobin levels of the umbilical cord vein compared to clamping the cord within 20 seconds. Thus, the study shows that early clamping (which is widely performed) is not justified.

The doctoral thesis Repercusiones clínicas y fisicoquímicas del tiempo de ligadura del cordón umbilical en recién nacidos a término (Clinical and Physiochemical Repercussions of Clamping Time of the Umbilical Cord of Newborns from Full-Term Pregnancies) analyzed a total of 151 umbilical cords of newborns from full-term pregnancies. In 79 cases, the umbilical cord was cut within 20 seconds, and in 72 cases it was clamped two minutes after the infant was expelled.

400+ News Categories

Popular News Topics

General News

Cancer News

Diet News

Drug News

Research News

Indian Health News

Alternative Medicine News

Corporate News

Education News

Health Conditions News

Lifestyle News **Dental News**

Sexual Health & Aging News

Women Health News Mental Health News

Diabetes News

Men's Health News **Hypertension News**

Organ Donation News

The study found that the partial pressure of oxygen in the umbilical artery of the newborn babies who had late clamping had risen, while there was a lesser need of oxygenotherapy after birth. There were no differences in the removal time of the placenta and the mother's bleeding after birth - one of the reasons why early clamping of the umbilical cord started to be practiced.

Page 1 Page 1 | 2 Next = Next News [™] **Child Health Related News** Eating Apples and Fish During Pregnancy Benefits Babies Where Indigenous Mothers Live Affects The Health Of Their Babies Baby's Birth Weight and Future Risk Has No Scientific Basis Second Hand Smoke may Prove Harmful for Underweight Babies Thymus Transplants Give Hope to Babies With Fatal Immune Disease Share this News: 💿 😭 📱 🥳 😉 🕻 🔭 📘 📝

Post Your Comments * Name : * Comment:

Ads by Google

Xavier Med Univ. **ValueMD**

US Clinicals, 100% Loan Programs Accredited MD Program in Aruba

www.valuemd.com/xavier

Advertise on this site

HEALTH VIDEOS



Ads by Google

Font Size

AAA

Is Your Heartburn Medici Working?




The Stress of Cancer: When Seek Help



and Its Cousir Heroin **Recognizing Opioid Abuse**







Last Updated - May 21, 2007 - Designed & Content Managed by Medindia Health Network Pvt Ltd. Hosted & Technical Support by Front Point Systems

Disclaimer - The contents of this site are for informational purposes only. Always seek the advice of a qualified physician for any doubts.

To Read full Disclaimer Click Here!

Advertise with us | Medindia Copyright | Privacy Policy | \odot All Rights Reserved 1997 - 2007