



*i*Village **Total I**

Articles Videos Books News **Images** Health & Medicine Space & Time

Mind & Brain

Plants & Animals

Earth & Climate

Share Blog Cite

Computers & Math Matter & Energy

Fossils & Ruins

Search

Science News

New Method Allows For The Early Diagnosis Of Pulmonary **Hypertension**

ScienceDaily (Nov. 14, 2007) — Researchers at the University of Granada (UGR) from the Systemic Autoimmune Diseases Unit and the Cardiology Service of the Hospital Clínico San Cecilio in Granada have developed an innovative system which will help doctors make the earliest diagnosis of pulmonary hypertension possible, which is the main cause of death for patients suffering from scleroderma, a rare disease which affects approximately 1,200 out of every million people.

See also:

Health & Medicine

- Hypertension
- Heart Disease
- Today's Healthcare · Diseases and Conditions
- Stroke Prevention
- Lung Disease

Reference

- Vein
- Emphysema
- Circulatory system
- Bronchitis

Until now, the only method to identify pulmonary hypertension was to perform a cardiac catheterization, an invasive technique consisting of the insertion of a hollow and flexible tube (a catheter) through the jugular vein directly into the right side of the heart. This technique is not advisable for all patients because of its complexity and risk.

The new technique was developed by doctor Lourdes López Pérez, and directed by lecturers Norberto Ortego Centeno and José Luis Callejas Rubio, from the Systemic Autoimmune Diseases Unit of the

Medicine Department, at the University of Granada.*

The great advantage of this system, which is based on the ultrasound technique, is that it allows specialists to measure the pulmonary artery systolic pressure so that pulmonary hypertension can be diagnosed earlier.

A destructive disease

Scleroderma is a systemic autoimmune disease, that is to say, it affects various organs of the body, and it is due to an alteration of the immune system. This disease is specially characterised by colour changing of the hands when exposed to cold, which is known as Raynaud's phenomenon, and causes skin swelling from fingers to wrists. Scleroderma is also characterised by damage in the skin of hands, neck and face. This damage resembles little spider veins known as telangiectasias. In addition to this, other organs such as the oesophagus, the kidneys or the lungs may be affected.

Lung damage is one of the most significant and serious symptoms, being the leading cause of death among patients suffering from scleroderma. The research team from the University of Granada maintained detailed medical records of each of the 41 patients included in order to accomplish this study. They also included data from both blood and respitatory tests. An echocardiogram was performed on all patients while they were resting and after physical effort on a special bicycle, called a bicycle ergometer.

From the results of this study, it has been concluded that there a group of patients who present a serious pathological increase of pulmonary artery systolic pressure when exercising. Therefore, these findings are very early evidence of scleroderma. Thanks to this early diagnosis a suitable treatment can be started to modify the scleroderma cycle, thus improving patient's prognosis significantly.

Early diagnosis

Given that pulmonary hypertension is a very serious complication of scleroderma, Lourdes López stresses the importance of diagnosing it as soon as possible in order to foresee the appearance of symptoms ascribed to it, such as breathing difficulty while partaking in physical activity or dyspnea.

She also states that using the bicycle ergometer to measure pulmonary artery systolic pressure of the patient after physical activity and performing an echocardiogram are very reliable methods, since they may help to identify scleroderma earlier.

The results of this new research confirms the importance of performing an echocardiogram on patients suffering from

Ads by Google

Advertise here

Stem Cell Stroke Therapy

Find out how your stroke can be treated with stem cells today!

www.xcell-center.com

COPD with Bronchitis?

There's a Proven Medication That May Help You **Breathe Easier**

www.gobreathe.com

Celsius Control Catheter

Inducing mild hypothermia during treatment for Cardiac Arrest/Stroke www.innercool.com

Coronary Artery Blockage

Method To Abate Coronary Artery Blockage www.coronaryarteryblockage.com

Metabolic Syndrome

Top world medical experts united to defuse the metabolic time bomb metabolic-syndrome-institute.org

Related Stories

Hormone Can Predict Pulmonary Hypertension And Potential Death Or Survival (Mar. 31, 2006)

 Measuring levels of a hormone called brain natriuretic peptide in individuals with serious lung disease can predict the presence of pulmonary hypertension and a patient's potential death or survival, ... > read more

Simple, Noninvasive Test Measures Survival

Time In Adult Pulmonary Hypertension (Nov. 3, 2006) — Researchers have developed a simple, noninvasive way to measure right ventricular function in the heart to predict survival of adults who suffer from pulmonary hypertension. The research appears in ... > read more

Heart Catheters Do Not Benefit Patients (Nov. 6,

2006) — Doctors should probably stop using pulmonary artery catheters because they do not benefit patients, said doctors from Australia recently in the British Medical ... > read more

New Approach To Pulmonary Hypertension Shows Promise (Jul. 9, 2007) — Researchers at

the University of Alberta have identified a "central command system" for pulmonary hypertension, a disease that currently has no cure and kills thousands each ... > read more

Sick Kids' Researchers Reverse Fatal

Pulmonary Hypertension (May 31, 2000) Research conducted at The Hospital for Sick Children (HSC) has opened up the possibility for a new treatment for pulmonary hypertension that may also be applicable to all types of blood vessel ...

14/11/2007 11:15 1 de 2

scleroderma, since this fatal complication may be indentified **Search ScienceDaily** before it becomes irreversible. Some of the results of this research have already been published in the European Journal of Echocardiography and Chest.

*They worked in collaboration with the cardiologists Eduardo Moreno Escobar and Pilar Martín de la Fuente.

Adapted from materials provided by Universidad de Granada.

Need to cite this story in your essay, paper, or report? Use one of the following formats: APA Universidad de Granada (2007, November 14). New Method Allows For The Early Diagnosis Of Pulmonary Hypertension. ScienceDaily. O MLA Retrieved November 14, 2007, from

/releases/2007/11/071113103804.htm

http://www.sciencedaily.com

Search Find with keyword(s):

the latest news stories, reference articles, science videos, images, and books. Ads by Google Advertise here

Enter a keyword or phrase to search ScienceDaily's archives for related news topics,

Levemir®

Information on insulin for healthcare professionals. www.NovoNordisk.com

Stroke

Answers to your questions about Stroke symptoms, treatments & more. health.ivillage.com

Scleroderma Treatments

Treatment Options, Latest Advances. Trustworthy, Current Report. www.scleroderma-info.com

Number of stories in

archives: 44,032

Email Newsletters RSS Newsfeeds

Feedback	we want to hear from you!
welcome both po	think of the new ScienceDaily we sitive and negative comments. ms using the site? Questions?
Your Name:	
Your Email:	
Comments:	
Click button to su	ıbmit feedback: Send It

Send It

About This Site | Editorial Staff | Awards & Reviews | Contribute News | Advertise With Us | Privacy Policy | Terms of Use Copyright © 1995-2007 ScienceDaily LLC — All rights reserved — Contact: editor@sciencedaily.com

14/11/2007 11:15 2 de 2