• rajesh on Benazir Bhutto killed in a sniper attack, five bullets were fired on her (Updated)

digg

- Bangkok will host Thailand's first ever BarCamp
- A Hypercar That Pays for Itself
- Benazir ignored warnings: Musharraf
- Britney Spears 5-hour date with paparazzo
- Search for aliens to get 500 times boost.
- Five technologies that may become a big hit in 2008
- Microsoft and Novell win Frost & Sullivan Award
- Green IT, XP reprieve and Linux among top 2008 IT forecasts
- Brit snaps UFO hovering over South-west England
- Pope setting up exorcism squads to fight rising Satanism

January 2008 MTWTFSS 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27

« Dec

Ads by Google

28 29 30 31

Pesticide Library Pesticide Summary Pesticides Health Pesticide What Is



100% of people have at least one type of pesticide in their body

January 6th, 2008 - 3:49 pm ICT by admin - Email This Post

Hesperia Granada

Disfruta de todas las ventajas de un hotel Hesperia desde 75 € www.hesperia.es

Washington, Jan 6 (ANI): Researchers at the Department of Radiology and Physical Medicine at the University of Granada, have discovered the presence of at least one kind of persistent organic compound (POC's) in almost 100 percent of Spaniards they examined.

It was also discovered that women and older people had a higher amount of pesticides, fungicides and insecticides present in the body, as compared to men and younger people respectively.

Vuelos desde Granada

Vuelos de línea Regular y Low Cost ¡Sólo 1€ de gastos de gestión! www.logitravel.com

The POCs are substances internationally classified as potentially harmful to ones health, and enter the human body through food, water or even air and get accumulated in the adipose tissue.

Ads by Google

The study led by Juan Pedro Arrebola Moreno, calculated the level of contamination of some POCs in people from two areas, an urban one and a semi-rural one, in order to discover factors linked to such levels: diet, lifestyle, activities or residence.

387 adults, from both sexes, volunteered for surgeries in hospitals taking part in the study. After their consent, the researchers took a sample of their adipose tissue in the surgery and later the volunteers were made to appear for a questionnaire about their place of residence, lifestyle, eating habits and activities throughout their life.

After analyzing the samples, the scientists measured 6 different POC concentration levels: DDE, a principal metabolite in DDT (a pesticide used in Spain until the 80's); hexachlorobenzene, a compound used as fungicide and currently released by industrial processes; PCBs: compounds related to industrial processes; and Hexaclorociclohexano, used as an insecticide and presently used in scabies and pediculosis treatment.

The researchers concluded that 100 percent of subjects analyzed had DDE in their bodies, a substance banned in Spain, and other very frequent components such as PCB-153 (present in 92 percent of people), HCB (91 percent), PCB-180 (90 percent), PCB-138 (86 percent) and HCH (84 percent).

Moreno explained that higher levels of toxic substances were detected in women compared to men and in older volunteers compared to younger people, possibly due to the great persistence of these substances in the environment, which results in their biomagnification in the food chain and in their bioaccumulation over time.

According to the scientists, another theory known as Efecto Cohorte (Cohort effect) may be behind high quantities of these substances in older people.

This theory indicated that people born in periods of higher contamination were more prone to suffer the consequences as compared to those born with the current bans on such pesticides.

Diet was also found to be an important factor in POC concentration, as the ingestion of some food materials, mainly of animal origin and high fat content, lead to a higher presence of these toxic substances in the human organism.

There are few studies in Spain measuring POC levels in wide samples of the population, which means that some compound levels in the general population are unknown, stated Moreno . (ANI)



Posted in Health Science |

- Related Stories
- Links for this article

Related Stories

- Organic food more nutritious than ordinary produce (November 14, 2007) 29% Match
- Fitness, not fat, may be stronger predictor of older adults longevity (December 5, 2007) 29% Match
- Married men have lower testosterone levels than bachelors (November 14, 2007) 27% Match
- Kids with Down syndrome more likely to have higher levels of obesity-related hormone (November 14, 2007) 25% Match
- Dramatic employment and poverty gap between fit people and disabled (November 25, 2007) 24% Match
- Vitamin D does not reduce overall cancer mortality (November 14, 2007) 23% Match
- Body abnormalities increase childhood cancer risk (January 2, 2008) 23% Match
- Maintaining zinc levels in blood may reduce pneumonia risk in nursing home elderly (November 14, 2007) 23% Match
- Researchers link 9/11 stress to low birth weight babies in New York (November 14, 2007) 22% Match
- Women with high BP at three-fold risk of developing diabetes (November 14, 2007) 22% Match

Categories

• RSS - Health Science

Tags

- RSS adipose tissue
- RSS concentration levels
- RSS eating habits
- RSS fungicide
- Ess fungicides
- RSS hexachlorobenzene
- RSS industrial processes
- RSS insecticide
- insecticides
- <u>RSS</u> <u>juan pedro</u>
- **RSS** <u>lifestyle activities</u>
- <u>metabolite</u>
- RSS organic compound
- RSS pcbs
- **PSS** pesticide
- RSS physical medicine
- RSS pocs
- RSS scabies
- spaniards
- RSS university of granada

