

ASSTEL

CosmosDirekt
127.100 €
für 5,18 € /Monat¹

Hannoversche Leben
129.570 €
für 5,18 € /Monat¹

150.000 €
für 5,18 € /Monat¹

innovationsreport

Forum für Wissenschaft, Industrie und Wirtschaft

Hauptsponsoren: SIEMENS n-tv Postbank

Datenbankrecherche:

Fachgebiet (optional):

GO

Home

Über uns

Media

English

FACHGEBIETE

SONDERTHEMEN

FORSCHUNG

B2B BEREICH

JOB & KARRIERE

SERVICE

NACHRICHTEN & BERICHTE

- Agrar- Forstwissenschaften
- Architektur Bauwesen
- Automotive
- Biowissenschaften Chemie
- Energie und Elektrotechnik
- Geowissenschaften
- Gesellschaftswissenschaften
- Informationstechnologie
- Interdisziplinäre Forschung
- Kommunikation Medien
- Maschinenbau
- Materialwissenschaften
- Medizintechnik**
- Medizin Gesundheit
- Ökologie Umwelt- Naturschutz
- Physik Astronomie
- Studien Analysen
- Verfahrenstechnologie
- Verkehr Logistik
- Wirtschaft Finanzen

Weitere Förderer



Google-Anzeigen

- Behandlung
- Therapie
- Studie
- Lexikon
- Medizin

Home → Fachgebiete → Medizintechnik → Nachricht

Design of a new system that improves hearing in patients who carry a cochlear implant

05.03.2008

The doctoral thesis of Isaac Manuel Álvarez Ruiz has been developed in the Departments of Department of Signal Theory, Telematics, and Communication, and Department of Surgery and its Specialties of the University of Granada, and directed by Professors Ángel de la Torre Vega and Manuel Sainz Quevedo.

Anzeige



The main aim of this work includes improving the acquisition of Compound Action Potential records, carrying out an application analysis in the cochlear implant programming and studying the distribution of refractory periods for the neuronal population that takes part in the generation of potentials. This doctoral thesis, carried out with the support of the Austrian company Med-El, is very useful for the manufacturers of cochlear implants as the results are a source of information for their present research lines.



Cochlear Implant Network

The definitive resource for hearing and balance disorders
www.cochlearnetwork.com

Hearing Loss

The New York Times has articles and information on the subject
www.nytimes.com

The cochlear implant is an electronic device designed to help patients with severe and deep hypoacusis who get very little or no benefit from conventional hearing aids. Patients who use them can detect environmental sounds, most of them are able to understand language with no need of lip-reading, and some of them can even use the phone.

This device is composed of an inner part, implanted surgically, and an outer part. In the inner part we mainly find the stimulus generator and an electrodes-bearing guide inserted all along the cochlea. The outer part is mainly composed of a microphone, a voice processor and a battery unit. So, the signal received by the microphone can be analyzed by the voice processor, which determines the instant in which electrodes should be activated as well as the level of the stimulus.

Specifically

The voice processor in the implant should be specifically programmed for each patient. In order to do so, a programmer performs some subjective tests on the patient. Carrying out this task can get especially hard with children or non-cooperative patients, whose interaction with the programmer is limited or void. So, the search for an objective measurement which provides the automatic programming for the cochlear implant is one of the main research lines opened by the manufacturers. The Compound Action Potential is included in this aim.

The author of this work, Isaac Manuel Álvarez Ruiz, states that "the major difficulty in the recording and processing of the Compound Action Potential is the stimulation device, insomuch that the stimulation signal interferes in time and frequency with the so-called potential that is meant to be observed". In his thesis, an algorithm has

B2B Suche

 GO

- ☒ Produkt / Dienstleistung
- ☐ Firma / Organisation

Aktuell

IronPort Threat Operation Center gibt Spam-Warnung für 2008

05.03.2008 | CeBIT 2008

Hotelmanagement: Intuition oder Controlling

05.03.2008 | Studien Analysen

Insulation monitors secure power supply

05.03.2008 | Maschinenbau



Veranstaltungen

Konferenz zu Erneuerbaren Energien in Washington

04.03.2008 | Veranstaltungsnachrichten

Elite und die Folgen - Die Auswirkungen der Exzellenzinitiative auf das Wissenschaftssystem

04.03.2008 | Veranstaltungsnachrichten

Internationale Bildungskonferenz am 6. und 7.3.



been developed to allow an evaluation of the quality of a Compound Action Potential recording. "This method lets us compare the most commonly used techniques in bibliography, as well as to develop new and improved techniques to reduce the stimulation device", points out the researcher.

The first task at the time of programming the cochlear implant is to decide which electrodes should be activated and subsequently, establish parameters for each one. So, this research has analyzed the relation between the Compound Action Potential and the cochlear implant voice processor programming map. "We have found that the appearance of the potential generally involves the electrode connection and not vice versa. According to the parameters of each electrode, it has been found that the information provided by these potentials is insufficient for an automatic adjustment of the processor, in comparison with the results of our researches. Our results prove that the Compound Action Potential allows the establishment of parameters of each electrode regarding the average values of the patient, with acceptable uncertainty levels". This information could be very useful as complementary information, or when the subjective information from the patient is void.

Refractory periods

In addition, a study has been conducted about the distribution of refractory periods for the neuronal population that takes part in the generation of this potential. In the average patient, the value of the refractory period of the fastest neuron is about 0.8ms and half of the neurons have a refractory period lower than 1.5ms. It has been found that, as the auditory experience of the patient increases, the refractory period of the neurons that takes part in it decreases down to a stable value, that is reached 3-4 months after the start up of the voice processor.

The results of this research have resulted in an article titled 'Generalized alternating stimulation: A novel method to reduce stimulus artifact in electrically-evoked compound action potentials', published in 2007 in the 'Journal of Neuroscience Methods'. Other articles are now under review.

Reference

Isaac Manuel Álvarez Ruiz. >Department of Signal Theory, Telematics, and Communication of the University of Granada.

Phone number: +34 626186403.

Email address: isamaru@ugr.es

Antonio Marín Ruiz | Quelle: alphagalileo

Weitere Informationen:

prensa.ugr.es/prensa/research/verNota/prensa.php?nota=508

Ihr Persönlichkeitstest

Detaillierte, wissenschaftliche Analyse Ihrer Persönlichkeit
www.personality100.de

Scheidenpilz behandeln

ohne Chemie und Antibiotika mit kolloidalem NanoSilizium
DynamSi
www.bnw-mineral.com/

Renditechance Agrarland

Investieren in landwirtschaftliche Flächen Osteuropas:
Gratis-Info
www.dikbg.com/agrarland

Webanalyse mit netmind

netmind: die Premium Webanalyse- Lösung für mehr
Umsatz
www.mindlab.de/index.php

in Darmstadt

04.03.2008 |

Veranstaltungsnachrichten

Live-Mitschnitte, Interviews und Hintergründe von den Meinungsführern aus Politik und Wirtschaft jetzt auf → www.euroforum.tv



Top

Artikel versenden

drucken

[Home](#) [Über Uns](#) [Partner](#) [Media](#) [Kontakt](#) [Sitemap](#) [find and help](#) [Englisch](#) [Impressum](#)

webdesign by freyhauer

CMS by Netzgut

Partnerseite: Xolopo

© 2000-2008 by innovations-report