



[News](#)
[Articles](#)
[Videos](#)
[Images](#)
[Books](#)
[Search](#)

[Health & Medicine](#)
[Mind & Brain](#)
[Plants & Animals](#)
[Earth & Climate](#)
[Space & Time](#)
[Matter & Energy](#)
[Computers & Math](#)
[Fossils & Ruins](#)

Science News

[Share](#)
[Blog](#)
[Cite](#)

[Print](#)
[Email](#)
[Bookmark](#)

New Geomorphological Index Created For Studying Active Tectonics Of Mountains

ScienceDaily (Sep. 16, 2008) — To build a hospital, nuclear power station or a large dam you need to know the possible earthquake risks of the terrain. Now, researchers from the Universities of Granada and Jaen, alongside scientists from the University of California (Santa Barbara, USA), have developed, based on relief data from the southern edge of the Sierra Nevada, a geomorphological index that analyses land form in relation to active tectonics, applicable to any mountain chain on the planet.

See also:

Earth & Climate

- [Natural Disasters](#)
- [Earthquakes](#)
- [Geology](#)
- [Tsunamis](#)
- [Volcanoes](#)
- [Renewable Energy](#)

Reference

- [Alpine Fault](#)
- [North Anatolian Fault](#)
- [Elastic-rebound theory of earthquakes](#)
- [Engineering geology](#)

Active tectonics comprise the most up-to-date deformation processes that affect the Earth's crust, resulting in earthquakes or recent deformations in the planet's faults and folds. This phenomena is analysed in geology research carried out before commencing engineering works.

Depending on the type of project (nuclear power stations or power stations, radioactive storage, natural gas or CO₂, large dams and tunnels, hydroelectricity projects...) and the type of earthquake (single or multiple), the time period for evaluating active tectonics varies between 10,000 and 100,000 years

for studies prior to beginning construction work.

The study, which is now published in the magazine *Geomorphology* and is the result of the doctoral thesis of Rachid El Hamdouni, Professor of the Department of Civil Engineering at the University of Granada, defines a new geomorphological index called Relative Active Tectonics Indexpos, which identifies four classes of active tectonics (from low to very high) and uses six geomorphological indicators.

"The main use of this new index is that it establishes a close relationship between this, the land forms, and direct evidence of active faults", El Hamdouni explained.

A seismic map for southern Spain

The indices are calculated with the help of Geographical Information Systems and teledetection programs in large areas which identify geomorphological anomalies possibly related to active tectonics. "This is really useful in southern Spain where studies on active tectonics are not very widely distributed", Chacón pointed out.

The study has focused on the Padul-Dúrcal fault and a series of associated fault structures on the edge of the Sierra Nevada, where over the last 30 years seismic activity has been recorded by the Observatory of the Andalusian Institute of Geophysics and Prevention of Seismic Disasters. Chacón explained that the map obtained with the new index depends exclusively on the land forms and divides the area studied into four parts, "of which two thirds of the total area is classed as having high or very high tectonic activity".

The Sierra Nevada is an Alpine mountain chain "with variable active tectonic gradients caused by the collision of Africa with Europe which has given rise to anticlines aligned from east to west, as well as the transverse extension with variable vertical gradients around 0.5 mm/year in normal faults", Chacón specified.

Ads by Google

[Advertise here](#)

5 Tips for a Flat Stomach

Stop making these 5 mistakes & you will finally lose your ugly belly.

[www.StomachFatsUgly.net](#)

Looking For Antibodies?

Find Quality Antibodies For Your Experiments. Order w/ Confidence!

[www.Millipore.com/Antibodies](#)

Empower Your Health

Sign Up Today and Get Access to Hot Topics Which Affect Your Health Now

[www.alsearsmd.com](#)

The Space Report

The premier analysis of the state of the space industry today

[www.thespacereport.org](#)

Support for FP7 proposal

Find partners and support for your FP7 proposal.

[www.temas.ch](#)

Related Stories



New Geomorphological Index Created For Studying Active Tectonics Of Mountains

(June 3, 2008) — To build a hospital, nuclear power station or a large dam you need to know the possible earthquake risks of the terrain. Now, researchers have developed, based on relief data from the southern edge ... [> read more](#)



Plan Developed For Restoring California Fisher Populations

(Jan. 15, 2008) — US Forest Service and U.C. Santa Barbara scientists believe they have identified the habitat needs for Pacific fishers, a rare California mammal that is a candidate for reintroduction efforts and ... [> read more](#)



Scientists Discover 'Moving Mountains'

(Aug. 11, 2004) — University of Nevada, Reno researchers have for the first time recorded a cluster of nearly 1,600 small earthquakes 20 miles beneath Lake Tahoe -- the world's second-largest alpine ... [> read more](#)

Geologists Study How Beachfront Property Turned Mountainous

(Sep. 7, 1999) — A modern cowboy would feel right at home along the Rocky Mountains in southern Wyoming, but if he could have been there 2 billion years ago, he would have needed a swimsuit—he'd be on the beach. ... [> read more](#)

Scientists Identified Earthquake Faults In Sichuan, China

(May 19, 2008) — Only last summer research published by earth scientists in the international journal *Tectonics* concluded that geological faults in the Sichuan Basin, China "are sufficiently long to sustain a strong ... [> read more](#)

Adapted from materials provided by [University of Granada - Communications Department](#).

Need to cite this story in your essay, paper, or report? Use one of the following formats:

- **APA** University of Granada - Communications Department (2008, September 16). New Geomorphological Index Created For Studying
- **MLA**