

- [Flight Status](#)
- [Jobs](#)
 - [Job Listings](#)
 - [Job Hunting Tips](#)
 - [In the Workplace](#)
 - [Career Growth](#)
 - [Changing Careers](#)
- [Cars](#)
 - [Buy](#)
 - [Sell](#)
 - [Weekly Specials](#)
 - [Research](#)
 - [drivingTV](#)
- [Homes](#)
 - [For Sale/Rent](#)
 - [Tips for Buyers & Sellers](#)
 - [Renovating](#)
 - [Decorating](#)
 - [Gardening](#)
 - [Condo Living](#)
 - [Vacation Homes](#)
 - [Green Homes](#)
- [Classifieds](#)
 - [Announcements](#)
 - [Dating](#)
 - [Obituaries](#)
 - [Job Listings](#)
 - [Car Listings](#)
 - [Real Estate For Sale/Rent](#)
 - [ShopLocal](#)
 - [Place an Ad](#)

Heavy pot smoking may trigger psychosis: Study

Reuters[Comments \(1\)](#)

- [Story](#)
- [Photos \(1 \)](#)



**Among individuals
who appear to be
"mentally well,"
heavy use of
marijuana may
predispose them to**

develop schizophrenia and other "psychoses," new research suggests.

**Photograph by: Christopher
Furlong, Getty Images**

NEW YORK (Reuters Health) - Among individuals who appear to be "mentally well," heavy use of marijuana may predispose them to develop schizophrenia and other "psychoses," new research suggests.

Among 92 patients, ages 18 to 65 years, who suffered a first episode of functional psychotic illness, more than half said they smoked marijuana daily or nearly every day and most of these individuals (66 percent) had no pre-existing signs of abnormal neurological development that would put them at risk for psychosis.

They had no family history of psychosis; they had been doing well academically; they had a group of friends and as such were not socially isolated; and they had good muscle coordination, Dr. Miguel Ruiz-Veguilla, of the Institute of Neurosciences of the University of Granada, Spain, and co-investigators explain in the latest issue of the journal Schizophrenia Research.

The remaining 43 percent of the study subjects with a first psychotic episode did have signs of abnormal neurological development that increased their risk of psychosis, such as a family history of psychosis and trouble in the social and academic worlds.

The new and most interesting study finding, wrote the researchers, is that mentally healthy adults who smoke marijuana daily or nearly every day may experience psychotic symptoms, while not having any of the risk factors for these symptoms.

This study, Veguilla further explained in comments to Reuters Health, suggests that patients with poor social and academic adjustment before the onset of illness "do not need an environmental factor in order to develop psychosis."

While people with good social and academic adjustment before the onset of psychosis illness - defined as have more than five friends; good academic function; good motor coordination and sensory integration; and no family history of psychosis -- need a strong environmental factor, such as smoking cannabis every day, to develop psychosis."

SOURCE: Schizophrenia Research 2009.

© Copyright (c) Reuters

- [E-mail this Article](#)
- [Print this Article](#)
- [Share this Article](#)

More on This Story

- [Study sheds lights on schizophrenia causes](#)
- [Recession depression](#)
- [Rules may be tightened on smoking medical marijuana](#)
- [Ont. wants federal clarity on medical pot](#)
- [Test kit highlights possible rise in prescription drug use among kids](#)
- [Marijuana may raise testicular cancer risk: Study](#)
- [Teen marijuana use no longer declining](#)

Story Tools

- [E-mail this Article](#)
- [Print this Article](#)
- [Comments \(1\)](#)
- [Share this Article](#)

Font:

- [*](#)