



LabAutomation
2010

Search Biosc

Search

 [RSS Feeds](#)  [Newsletters](#)  [Bookmarks](#)

[\[-\] Text](#) [\[+\]](#)

Gene increases effectiveness of drugs used to fight cancer and allows reduction in dosage

Featured In: [Disease Research](#)

By EurekAlert

Tuesday, November 24, 2009

Researchers at the [University of Granada](#) have found a suicide gene, called 'gene E', which leads to the death of tumour cells derived from breast, lung and colon cancer, and prevents their growth. The importance of this new gene is that its use to fight cancer can reduce the potent drugs that are currently used, so that could mean more effective treatment for cancer.

This research was conducted by **Ana Rosa Rama Ballesteros**, from the Department of Anatomy and Human Embryology at the [University of Granada](#), and directed by professors **Antonia Aránega Jiménez**, **José Carlos Prados Salazar** and **Consolación Melguizo Alonso**. Its aim was to study the possibility of reducing the dosage of drugs currently administered to cancer patients using combination therapy with suicide gene E.

Scientists from the UGR have shown that the bacteriophage phiX174 killer gene called E, can be used to induce death in tumour cells. So far, attempts to use many chemotherapeutic (cytotoxic) agents similar to the E gene have shown severe limitations resulting from their toxicity and their poor affinity with the tumour.

Advantages of gene therapy

As Ana Rosa Rama explains, chemotherapy, radiotherapy and surgery show at present "limited" results in advanced stages of cancer. "That is why it is urgent to find new therapies, and gene therapy has emerged as a potentially powerful therapeutic platform." Her work has shown that "it is possible to use gene therapy as an aid to chemotherapy, improving its results when it comes to attacking cancer, thus allowing the dosage of agents to be reduced and contributing to a reduction in side effects for the patient."

In order to understand how the E gene works, the researchers conducted studies using various techniques. The results indicate that the E gene's mechanism of action is to induce apoptosis (cell death), possibly through mitochondrial injury.

Therefore, they stress that "this new E gene appears as an ideal candidate to be transfected into tumour cells in order to induce apoptosis, possibly through mitochondrial activation, and to increase the sensitivity of these cells to the action of the drug developed specifically to act on them."

The results of this research suggest the possibility of reducing the concentration of chemotherapeutic agents in current use with cancer patients. Thus, in lung cancer cell line A-549, scientists from the UGR achieved a 14% inhibition of tumour growth and reduced by 100 times the dose of Paclitaxel agent when it was combined with gene E. In the case of colon cancer, the results were similar. However, the most relevant fact was found in the breast cancer cell line MCF-7, in which the dose of the chemotherapeutic agent, doxorubicin, was reduced by 100 times, reaching up to a 21% greater inhibition of tumour proliferation when combined with gene E. Currently, researchers from the UGR are in the process of obtaining a patent for gene E.

SOURCE

Join the Discussion

Rate Article:  Average 0 out of 5
[register](#) or [log in](#) to comment on this article!

0 COMMENTS
ADD COMMENT

   

 Email  Print

 1 tweet
 retweet

News

Supplier Update

- Elite Del. antique car expert seeking apprentices
1 hour ago
- INSIDE WASHINGTON: Stream of WH health care visits
1 hour ago
- BECK for November 25, 2009 - Part 1
7 hours ago
- Onex raises US\$151M in sale of part of its stake Emergency Medical Services
9 hours ago
- La Jolla Pharma axes plan to dissolve company
11 hours ago
- FDA grants full approval to J&J's HIV drug
12 hours ago
- Implant-based cancer vaccine is first to eliminate tumors in mice
13 hours ago

[more NEWS >>](#)

LABORATORY EQUIPMENT
Webcast Series

NOW ON DEMAND

Analytical Techniques for
Monitoring Bioethanol

Sponsored by:

 **PerkinElmer**
For the Better

 **phenomenex**
...breaking with tradition

REGISTER FREE

Most Popular

Related Content

- Cellular Source of Most Common Type of Abnormal Heart Beat Found
Nov 21
- Higher Carotid Arterial Stenting Rates Associated with Poorer Clinical Outcomes
Nov 21

Promoting healthy skepticism in the news:
Helping journalists get it right