

redOrbit

[LOGIN](#)  
[SIGN UP](#)  
[EMAIL](#)  
[SUGGESTIONS](#)

[HOME](#) [COMMUNITY](#) [NEWS](#) [VIDEO](#) [IMAGES](#) [SPACE](#) [SCIENCE](#) [TECH](#) [HEALTH](#) [EDUCATION](#) [TOPICS](#) [SHOP](#) [SITEMAP](#)

SEARCH

[Space](#) [Science](#) [Technology](#) [Health](#) [General](#) [Sci-fi & Gaming](#) [Oddities](#) [International](#) [Business](#) [Politics](#) [Education](#) [Entertainment](#) [Sports](#)
[Print](#) [Comment](#) [Font Size](#) [Digg](#) [del.icio.us](#) [Discuss article](#) [Buzz up!](#) [0](#) [Me gusta](#)

## A Protein Called CFLIP Makes Tumor Cells In Breast Cancer Resistant To Treatments

Posted on: Tuesday, 14 December 2010, 20:23 CST

Researchers at the Andalusian Institute for Molecular Biology and Regenerative Medicine (CABIMER) and the University of Granada found that cFLIP –an inhibitor of death ligand-induced apoptosis– is not only essential in breast tumor cells resistance to TRAIL treatments (a death ligand with a potent therapeutic potential against cancer), but this protein is also key to the survival of such cancer cells.

Researchers proved that a variation in the expression of this protein may lead to the normal development of breast epithelium. This is an important finding to be considered in the design of cFLIP-targeted therapies against cancer.

The research conducted by Rosario Yerbes Cadenas, PhD candidate at the University of Granada, was led by professor Abelardo López Rivas, of CABIMER, and was aimed at analysing the potential of cFLIP inhibitors in cancer therapies.

At present, TRAIL is a death-ligand of the TNF family, with significant therapeutic potential against cancer, basically due to its ability to induce apoptosis in cancer cells without displaying significant toxicity toward normal cells. However, there are tumor cells that are resistant to TRAIL-induced apoptosis for unknown causes.

### A Key Component

This study analysed the role of cFLIP in breast cancer cells' resistance to TRAIL-induced apoptosis. Thus, researchers concluded that cFLIP is key in these cells' resistance to TRAIL. Such conclusion was drawn from the evidence that the inhibition of their expression through treatments with Doxorubicin (anthracycline, widely used in chemotherapy) or with SAHA (Histone deacetylases inhibitor), as well as the silencing of its expression through cFLIP siRNA oligos (small interfering RNA), resulted in the sensitisation of breast cancer cells to TRAIL-induced apoptosis.

The authors of this research proved that cFLIP plays a survival role in tumorous and non-tumorous breast epithelial cells, since the inhibition of its expression induces apoptosis. This type of apoptosis requires the formation of the death-inducing signalling complex, which includes TRAIL-R2 receptor, adapter molecule FADD and procaspase-8- but is TRAIL-independent itself.

Conversely, in the light of the cFLIP relevance in controlling apoptosis, researchers studied the role of cFLIP in breast epithelial cells MCF-10A morphogenesis -a process where apoptosis plays an essential role. Thus, cFLIP/cFLIPS overexpression inhibits lumen formation in acini from breast epithelial cells when they are cultured in a 3D extracellular matrix (3D cultures). Additionally, inhibition of cFLIP expression prevents the development of acini, since cells with low expression of cFLIP are unfeasible.

For this reason, regulation of cFLIP expression was very relevant to this research. Scientists determined that the PI3K/AKT signalling pathway is not the main responsible for cFLIP synthesis in breast cancer cell, but may be it is NF-κB pathway.

Additionally, this study revealed that the ubiquitin-proteasome system plays a key role in cFLIP cell degradation. At present, researchers are trying to identify E3-ubiquitin ligase protein, responsible for cFLIP degradation by such system.

---

On the Net:

[University of Granada](#)

[More News in this Category](#)

### Related Articles

[VCU Massey Discovery Could Lead To Breakthrough For Non-Small Cell Lung Cancer](#)  
[Patient Enrollment Complete in Tragara Pharmaceuticals' Phase II Trial of Capoxigem\(R\) in Non-Small Cell Lung Cancer](#)  
[Regulatory Role Of Key Molecule Discovered](#)  
[New Pfizer Data Presented on CP-751,871 Across Non-Small Cell Lung Cancer and Ewing's Sarcoma](#)  
[Physician Insights on Key Attributes - Non-Small-Cell Lung Cancer](#)  
[Preclinical Data Show Alfacell's ONCONASE\(R\) Inhibits Tumor Growth of Non-Small Cell Lung Cancer, Breast Cancer Cells](#)  
[View Scientific Rationale for Novel Therapeutics in Lung Cancer, And the Results of Clinical Trials to Date Inside 'Triple Analysis: Non Small Cell Lung Cancer, Protein Kinase Inhibitors & Anti-Angiogenic Agents'](#)  
[FoldRx Announces Study Defining Role of Key Protein and Biological Pathway Implicated in Parkinson's Disease; Collaborators' Research in Science Points Way to New Treatment Approaches and Validates Company's Discovery Engine](#)  
[Loss of PNA Staining in Mouse Aortico-Pulmonary Septum is Associated With Mesenchymal Cell Apoptosis](#)  
[The Role of Whole-Body Bone Scanning and Clinical Factors in Detecting Bone Metastases in Patients With Non-Small Cell Lung Cancer\\*](#)



Smarter City

Dec 14, 2010, 9:58 am

[Solar Corona Observed by SDO](#)

Dec 14, 2010, 9:55 am

[Texting Examined for 911 Emergencies](#)

Dec 14, 2010, 9:45 am

[New Technology Saves Distracted Drivers](#)

Dec 14, 2010, 9:39 am

[Clean Energy: OpenHydro Tidal Technology](#)

Dec 14, 2010, 9:22 am

[Urban Heat Islands](#)

Dec 14, 2010, 9:17 am

[IBM Helps Corpus Christi Build a Smarter City](#)

[More Videos](#)

### Sponsors

[Breast Cancer Awareness Month](#)

[Cancer Symptoms](#)

[Advanced Breast Cancer](#)

[Cancer Information](#)

[Find Cancer](#)

[Cancer Research](#)

[Breast Cancer Survivors](#)

[Breast Cancer Information](#)

[About Cancer](#)

[Comprehensive Cancer Treatment](#)

[Cancer Treatment](#)

### Related Videos

[New Cancer Treatment Uses Cancer-resistant Blood](#)  
[High-Tech Molecular Breast Imaging](#)