



# DailyIndia.com

Your daily news source for India and more

 Search

[Home](#) [National](#) [World](#) [Business](#) [Sports](#) [Cricket](#) [Entertainment](#) [Health](#) [Science](#) [Bollywood](#) [Celebs](#) [Geekwerks](#)

## Protein that makes tumor cells in breast cancer resistant to treatments

From ANI

Washington, Dec 15: Scientists have found a protein that makes tumor cells in breast cancer resistant to treatments.

### Breast Cancer Trials

Learn about phase III trials for metastatic breast cancer disease  
[www.thewideprogram.com](http://www.thewideprogram.com)

### Interventional Oncology

Plan. Guide. Assess. Find out more at GE Healthcare.  
[www.gehealthcare.com/oncology](http://www.gehealthcare.com/oncology)

### X-tremeGENE Transfection

New efficient Roche Reagents – Request your free sample today!  
[www.roche-applied-science.com](http://www.roche-applied-science.com)

### Hyperthermia in Frankfurt

Extreme heat damages cancer cells A gentle biological cancer therapy  
[www.hyperthermie-zentrum.de](http://www.hyperthermie-zentrum.de)

Ads by Google

Researchers at the Andalusian Institute for Molecular Biology and Regenerative Medicine (CABIMER) and the University of Granada found that cFLIP -an inhibitor of death ligand-induced apoptosis- is not only essential in breast tumor cells resistance to TRAIL treatments (a death ligand with a potent therapeutic potential against cancer), but this protein is also key to the survival of such cancer cells.

They found that a variation in the expression of this protein may lead to the normal development of breast epithelium.

The study analyzed the role of cFLIP in breast cancer cells' resistance to TRAIL-induced apoptosis. Thus, researchers concluded that cFLIP is key in these cells' resistance to TRAIL.

Such conclusion was drawn from the evidence that the inhibition of their expression through treatments with Doxorubicin (anthracycline, widely used in chemotherapy) or with SAHA (Histone deacetylases inhibitor), as well as the silencing of its expression through cFLIP siRNA oligos (small interfering RNA), resulted in the sensitisation of breast cancer cells to TRAIL-induced apoptosis.

The researchers have proved that cFLIP plays a survival role in tumorous and non-tumorous breast epithelial cells, since the inhibition of its expression induces apoptosis.

Copyright Asian News International/DailyIndia.com

Currently trending: [Cricket](#), [Shane Warne](#), [Angelina Jolie](#), [Lindsay Lohan](#), [Oprah Winfrey](#), [Michael Clarke](#), [Liz Hurley](#), [Shahid Afridi](#), [Lady Gaga](#), [Miley Cyrus](#)

Share it:



[Related News](#)

[Most Popular Stories](#)

[Related News](#)

- [A quarter of Brit kids are overweight or obese by the age of 5](#)
- [New discovery shows promise against pancreatic cancer](#)
- [Bacteria that live on and in humans orchestrate health and disease](#)
- [How 'chameleon-like' HIV virus eludes treatment by mutating](#)
- [Southampton Varsity probing if deaf have better language skills](#)
- [3D TVs 'can cause headaches, nausea'](#)

[Most Popular Stories](#)

1. [Flat-dwelling kids 'more likely to be exposed to tobacco smoke'](#)
2. ['Miracle' blood transplant 'cures American man of HIV'](#)
3. [Justin Bieber, Inception, Gulf oil spill are this year's top Twitter Trends](#)
4. [Oprah Winfrey stuns Sydney audience with bountiful gifts](#)
5. ['Famine survival gene' offers clues to cause of obesity](#)