Home » General, Health Care

# Research: Eating disorders more prevalent among Muslim adolescents than Christian

Written By: ameilia on March 9, 2011 0

The incidence of eating disorders was found to be 2.3-fold higher among Muslim adolescents than among their Christian classmates. Similarly, body dissatisfaction was 1.8-fold higher in the former group. Finally, as a general conclusion, an average of one in four adolescents suffers some type of eating disorder, and 15% suffers body dissatisfaction. These were the conclusions drawn of a research conducted at the University of Granada.

The sample was taken from a Spanish multicultural city, Ceuta, where different religious groups have co-existed for centuries -Christian, Muslim, Jews-, and where about 30% of the total population is Muslim. The goal of this study was to assess the incidence of eating disorders and body dissatisfaction in a non-clinical sample of adolescents belonging to different religious groups, and assess the relationship between religious affiliation and health disorders.

To carry out this research, the authors examined a sample of 493 students aged between 12 and 20 years from three public schools in Ceuta, including secondary school.

### Pioneer Study

The author of this study, Francisco Javier Ram-n Jarne, belongs to the Institute of Neuroscience "Federico Ol-riz" of the University of Granada. This research "addresses a subject that has never been studied in Spain before: the relationship between religious affiliation, eating disorders and body image perception. As regards other international studies, this is one of the few comparative studies ever made by religious affiliation (Christian and Muslim) and gender". So far, only one similar study had been developed in Israel with a sample of Christian, Muslim and Jew adolescent girls.

The recent massive migratory movements -mainly from countries of the Maghrib-experienced in Spain have transformed the religious affiliation of its population. "This makes the results of this study very interesting from the point of view of preventive medicine and public health", Francisco Javier Ram-n Jarne states.

The University of Granada researcher pointed out that "affiliation to the Islamic religion taken as a socio-cultural factor can also be associated to these disorders, even although the thin ideal is not associated to the Islamic culture".

Source: University of Granada

## Add this article to the following Bookmark Services

Digg this! Add to del.icio.us! Stumble this! Add to Techorati! Share on Facebook! Seed Newsvine! Reddit! Add to Yahoo!

### Have your say:

You must be <u>logged in</u> to add your Say.

# More about this from BIOMEDME.COM

Study highlights | Only half of | New Specialist | T | lack of treatment | for adolescents | affected with | severely impairing | Entire New Specialist | T | Library offers | E | E | treatment for | C | C | treatment for | Severely impairing | Entire New Specialist | T | Entire New

disorders
A recent study by
Merikangas and
colleagues published in
the January 2011 issue
of the Journal of the...

severely impairing mental disorders receive treatment A recent study by Merikangas and colleagues published in the January 2011 issue of the Journal of the... New Specialist
Library offers
treatment for
eating disorders
Eating Disorder Hope, a
respected independent
resource for individuals
suffering from anorexia,
bulimia,...

The Impact Of
Eating Disorders
On NativeAmericans
Revealed By New
Study
Scientists in Connecticut
have carried out one of
the first psychological
studies into eating

disorders..

Registered Users

Username

Password

LOGIN Premember Register
Recover password

Linked in facebook

### MAIN MENU

- Biomed Articles
- Our Services
- About Middle East
- Biomed Events
- <u>Useful Links</u>
- Country Profiles
- <u>Downloads</u> <u>Biomed</u>
- Professionals Interviews
- <u>Videos</u>
- Membership Courses we offer
- Contact us
- Privacy Policy
- Terms & Conditions
- <u>Submit</u> <u>Pressrelease</u>
- <u>Biomed</u> <u>Professions</u>

2 de 3