

1 de 4 06/04/2011 11:07

Ads by Google
Bipolar Type 2
Social Work
Social Anxiety

Alcoholism is specially conflictive at family level, and alcohol is the most abused substance in patients with bipolar disorder. "Detecting alcohol abuse in patients is specially important to improve family relations".

The results obtained in this study were recently presented in the IV Portugal-Spain Meeting for Therapeutical Adherence held in Oporto, Portugal, where it was awarded the third research prize.

Sources: University of Granada, AlphaGalileo Foundation.

Please rate this article: Patient / Public: or Health Professional: (Hover over the stars then click to rate)





Add Your Opinion

Please note that **we publish your name**, but we **do not publish** your email address. It is only used to let you know when your message is published. We do not use it for any other purpose. Please see our privacy policy for more information.

If you write about specific medications or operations, please **do not** name health care professionals by name.


All opinions are moderated before being included (to stop spam)

Your Name:* E-mail Address:* Title For Opinion:* Opinion:*
This is to help prevent SPAM submissions. Please enter the words exactly as they appear, including capital letters and punctuation.* Type the two words:
Submit! * Fields marked with a * need to be filled in before you hit the submit button.

Contact Our News Editors

For any corrections of factual information, or to contact the editors please use our feedback form.

Please send any medical news or health news press releases to: pressrelease@medicalnewstoday.com



2 de 4 06/04/2011 11:07