

Print Comment Font Size Digg del.icio.us Discuss article Buzz up! 0 Me gusta

Half The Patients With Bipolar Disorder Suffers Work, Social Or Family Disabilities

Posted on: Tuesday, 5 April 2011, 15:56 CDT

Such was the conclusion obtained in a study conducted at the University of Granada that was recently awarded a prize in the IV Spain-Portugal Meeting on Therapeutical Adherence held in Oporto (Portugal). This research study analyzed the factors associated to higher work, social and family disabilities in a sample of 108 patients suffering from bipolar disorder.

According to the study, 50% of patients suffering from bipolar disorder suffers some type of work, social and family disability, and approximately 20% present some disorder at the three levels.

This was the conclusion drawn in a scientific article recently published in the prestigious journal Psychiatry Research, prepared by Dr. Luis Gutiérrez Rojas, a member of the Research Group of Psychiatry Research and Neuroscience of the University of Granada coordinated by professor Manuel Gurpegui Fernández de Legaria. This study analyzed the factors associated to serious work, social and family disabilities in a sample of 108 patients suffering from bipolar disorder.

This research revealed that work disability –that is, difficulty to perform normal job duties– in these patients was associated to high recurrence of manic episodes, as well as to recurrent psychiatric hospitalization –high-intensity episodes–, depression and low educational levels. Furthermore, nicotinic dependency –strong addiction to tobacco– can be more disabling in patients with bipolar disorder than depression.

Social Support

Gutiérrez Rojas states that social disability –difficulty to establish relations out of the family and to achieve social integration– in these patients is associated to higher hospitalization rates, episodes of depression and active depression symptoms. "Receiving social support is associated to lower social disability in these patients" –the author states.

Lastly, family disabilities –difficulty to have good relations with family members– was found to be associated to hospitalization, manic episodes, symptoms of depression and to higher scores in the CAGE scale, which measures addiction to alcohol.

The University of Granada researcher adds that, to avoid disability in patients with bipolar disorder "recurrence of depressive and manic episodes must be avoided, and physicians should treat these episodes promptly to avoid hospitalization".

Gutiérrez Rojas states that dependency to tobacco –apart from serious physical sequels– is associated to a worse bipolar disorder prognosis. Simultaneously, social support should be improved or sought for patients deprived of it. For instance, home assistance –instead of financial support– can be sought within the framework of the Spanish Law of Dependency. Patients can also be advised to attend patient meetings in search of information and support".

Alcoholism is specially conflictive at family level, and alcohol is the most abused substance in patients with bipolar disorder. "Detecting alcohol abuse in patients is specially important to improve family relations".

The results obtained in this study were recently presented in the IV Portugal-Spain Meeting for Therapeutical Adherence held in Oporto, Portugal, where it was awarded the third research prize.

On the Net:

[UGR](#)
[Psychiatry Research](#)

[More News in this Category](#)

Related Articles

- [American Airlines and American Association of People With Disabilities Launch Inaugural 'Altitude Award' Call for Entries](#)
- [Home Oxygen Patients and Family Members Visit Capitol Hill to Ask Congress to Protect Medicare Home Oxygen Therapy Benefit](#)
- [American Airlines and American Association of People With Disabilities Team up to Honor Best TV Commercial](#)
- [Study Finds that Dying Patients and Families Suffer from Lack of Continuity of Physician Care](#)
- [Alexza Announces Preliminary Results From Its AZ-002 \(Staccato\(R\) Alprazolam\) Phase 2a Proof-of-Concept Trial in Patients With Panic Disorder](#)
- [The Patient Recruiting Agency\(TM\) Announces New Study of Major Depressive Disorder](#)
- [National Eating Disorders Association Reacts to 'Guidelines' Announced Today By Council of Fashion Designers of America](#)
- [Avera Pharmaceuticals Completes Enrollment in Phase 2 Social Anxiety Disorder Study](#)
- [New Study Reports Venlafaxine Extended Release May Help Adult Patients Prevent New Episodes of Depression for Up to Two Years](#)
- [Randomized, Double-Blind, Placebo-Controlled Trial of Fluoxetine Treatment for Elderly Patients With Dysthymic Disorder](#)



Stay Connected

Discover the best free real-time news, networking and information portal on the web... [Learn more](#)



Create Your Own News!

NewsCircles are a quick, convenient way to create and publish your own customized news portals... [Learn more](#)



Never Miss A Beat

Get the most up to date breaking new stories as they happen across the globe...



Top Industry News

Choose an industry to view real-time news clustered and organized from over 25,000



Apr 5, 2011, 9:48 am

[Gardening Has Many Benefits](#)

Apr 5, 2011, 9:46 am

[Plastic Surgery is on the Rise Among Men](#)

Apr 5, 2011, 9:35 am

[Lord of the Tree Rings](#)

Apr 5, 2011, 9:21 am

[Expedition 27 Launch](#)

Apr 5, 2011, 9:12 am

[Cyberchondria](#)

Apr 5, 2011, 9:07 am

[Kids Should Remain in Rear-Facing Car Seats Until 2](#)

[More Videos](#)

Sponsors

[Read On...](#)

Advertisement

Related Videos

- [New Options for Treating Bipolar Disorder](#)
- [Depression Screening May Not Benefit Heart Patients](#)
- [PANDAS](#)
- [Lack of Sunlight Leads to Depression](#)
- [Scientists Developing Paralysis Cure](#)
- [Researchers: Marijuana May Help MS Patients](#)