



Half of All Bipolar Disorder Patients Grapple With Work, Social or Family Disabilities

by Tanya Thomas on April 10, 2011 at 10:17 AM

Adolescent Psychiatry

Official Journal - American Society of Adolescent Psychiatry
www.benthamscience.com/aps

Ask a Psychiatrist Online

5 Psychiatrists Are Online. Current Wait Time: 7 Minutes.
Health.JustAnswer.com/Psychiatry

Private Health Insurance

Different levels of cover available
 Affordable policies in English
www.nashwarren.co.uk

 Select Language

Gadgets powered by Google



Ads by Google

Mental Health News

[News](#) [Comments](#) 7 retweet Text

www.lumosity.com
 Ads by Google


A study conducted at the University of Granada has found that 50 pc of patients with bipolar disorder suffer some type of work, social and family disability, and approximately 20 pc present some disorder at the three levels.

The research study analyzed the factors associated to higher work, social and family disabilities in a sample of 108 patients suffering from bipolar disorder.

Latest News Most Popular Most Commented

- Mind Reading Computer Technology
- New Method to Turn Blood Cells into Beating Heart Cells
- Simple Way to Boost Your Brain - Learn Like Baby
- Freeway Air Pollution Boosts Brain Damage in Mice
- New Insight into Acute Pancreatitis

News Quick Links Special Reports

- News Central
- Latest Health News
- News Category (500+)
- Popular News
- Health News and Press Release
- News Photo Gallery
- News Video Gallery
- News From Other Resources
- Health Watch
- Health In Focus
- Breaking Health News
- Celebrating Life
- Medindia - Exclusive
- India Special
- Lifestyle and Wellness

News Archive

Date :

Category :

Keyword :

Select a News Category

- Kate Middleton's Unpleasant Experience at School [Read More](#)
- Bhagwan Satya Saibaba's Condition is Stable Now [Read More](#)
- Andrew Lloyd-Webber Rendered Impotent After Prostate Surgery [Read More](#)

[More Celebrity News](#)

This research revealed that work disability -that is, difficulty to perform normal job duties- in these patients was associated to high recurrence of maniac episodes, as well as to recurrent psychiatric hospitalization -high-intensity episodes-, depression and low educational levels. Furthermore, nicotinic dependency -strong addiction to tobacco- can be more disabling in patients with bipolar disorder than depression.

Luis Guti rrez Rojas, a member of the Research Group of Psychiatry Research and Neuroscience of the University of Granada states that social disability -difficulty to establish relations out of the family and to achieve social integration- in these patients is associated to higher hospitalization rates, episodes of depression and active depression symptoms.

"Receiving social support is associated to lower social disability in these patients," said the

author.

The University of Granada researcher added that, to avoid disability in patients with bipolar disorder "recurrence of depressive and maniac episodes must be avoided, and physicians should treat these episodes promptly to avoid hospitalization".

Rojas stated that dependency to tobacco -apart from serious physical sequels- is associated to a worse bipolar disorder prognosis. Simultaneously, social support should be improved or sought for patients deprived of it.

Adapted by Google from: [PBA University](#), [PBA University](#), [Medical Diet](#), [Tours, Granada](#)

Page 1

Page 1 | 2

Next ➔

PREVIOUS STORY

[Genetic Breakthrough to Major Cause of Kidney Disease](#)

NEXT STORY

[Accelerated Immune Cell Aging Linked to Chronic Depression](#)

[Email](#) [RSS Feeds](#) [Print](#) [Save](#) [Link](#) [Syndicate](#) [Comments](#) [SHARE](#) [Facebook](#) [Twitter](#) [LinkedIn](#)

More News on: [Amoebic Dysentery](#), [Quiz on Depression](#), [Sleep Problems in Children](#)

Comment & Contribute

Be the first to comment!

* Your comment can be maximum of 2500 characters

- ☒ Notify me when reply is posted
☒ I agree to the [terms and conditions](#)

[Post Comments](#)


Medindia en Facebook

Me gusta

A 3,513 personas les gusta **Medindia**.













Are you a member of Facebook? [Become a member of Medindia](#) and get breaking news delivered to your Facebook page.

Medindia Health Network

[Support Groups](#) [Members](#) [Discussions](#)



Posted By : [Mahadi](#) on Thu, Apr 7 2011

[Dear sir, Hi! how are you. sir some advise needed, I am mahadi, 29 years, my Amo...](#)

Under : [Amoebic Dysentery / Amoebiasis](#) Support Group : [Amoebic Dysentery / Amoebiasis](#)

No. of Posts: 23 Views: 1346 [Reply](#)



Posted By : [rknaveen](#) on Wed, Oct 20 2010

[I too had same kind of recurring problem while taking allopathic medicene. But I...](#)

Under : [Amoebic Dysentery / Amoebiasis](#) Support Group : [Amoebic Dysentery / Amoebiasis](#)

No. of Posts: 23 Views: 1346 [Reply](#)



Posted By : [purpleenergy](#) on Tue, Sep 7 2010

[Tesla Plates are ideal for childrens and pregnant mothers. It is a natural pai...](#)

Under : [Lack of Sleep May Put Young Kids At Risk of Obesity](#) Support Group : [Sleep Disorder:Sleep](#)

[Problems in Children](#) No. of Posts: 2 Views: 675 [Reply](#)

Total Members - 38683
Support Groups - 445

[Login](#)


[Join Now >>](#)



Mental Health Related News



Accelerated Immune Cell Aging Linked to Chronic Depression



Online Mental Health Modules Reduce Anxiety or Depression



Greater Control Over Thoughts Possible With Real-Time Brain Feedback



Memory Suffers Due to Stress

[Read More](#)