

Consuming probiotics for a month helps diminish fat accumulation in the liver, new study says

Public Release: 18-Jul-2014 Consuming probiotics for a month helps diminish fat accumulation in the liver, new study says

Scientists at the University of Granada have made important strides in the fight against the Non-Alcoholic Fatty Liver Disease, which is closely related to obesity and diabetes

University of Granada

[Print E-Mail](#)

Spanish scientists have demonstrated through an experiment on obese rats that the consumption of probiotics during thirty days helps diminish the accumulation of fat in the liver. This new finding, published today by the journal *PLOS ONE*, is a great step forward on the fight against the Non-Alcoholic Fatty Liver Disease (NAFLD), which is closely related to obesity and diabetes.

Researchers from the 'Nutrition Biochemistry: Therapeutic Applications' group (CTS-461) and the José Mataix Institute for Nutrition and Food Technology at the University of Granada have demonstrated that the administration of three probiotic strains diminishes the accumulation of fat in the liver of obese rats.

The accumulation of fat in the liver is called steatosis and it constitutes the first stage in the NAFLD disease, which is closely related to obesity and diabetes. Given that the prevalence of these two pathologies does not cease to increase, NAFLD has also become a health problem that affects millions of people throughout the world.

Living or dead microorganisms

Probiotics are microorganisms (bacteria or yeasts) with healthy effects upon individuals that consume them in adequate doses. They were traditionally considered to be living microorganisms, but the concept was widened since some dead microorganisms, or even their components, can display probiotic properties.

University of Granada researchers worked with three strains which are custodied at the Collection Nationale de Cultures de Microorganismes (CNCM) of the Pasteur Institute: *Lactobacillus paracasei* CNCM I-4034, *Bifidobacterium breve* CNCM I-4035 and *Lactobacillus rhamnosus* CNCM I-4036. During their first experiment, conducted on healthy volunteers, researchers demonstrated that all three of them are perfectly tolerable and safe for human consumption.

In this current study, the strains were administered during thirty days in the diet of Zucker rats. These rats develop obesity due to a mutation in the gene that codifies the receptor or leptine, a hormone that transmits a sensation of satiety to the organism. Zucker rats are among the best characterized genetic models.

In their article, the authors describe that the administration of probiotics led to an accumulation of lipids (most of them triacylglycerides) in the liver which was significantly lower than that occurring in rats fed with a placebo.

"This new finding went hand in hand with lower values in proinflammatory molecules (tumor-necrosis factor, interleukin 6 and liposaccharide) in the serum of rats fed with probiotics. These effects were not observed in those

According to these researchers, this liver disease will not be cured with probiotics, but these microorganisms can certainly be used as support therapy in joint use with other treatment.

###

This study has been financed by the private company HERO SPAIN S.A.

Disclaimer: AAAS and EurekAlert! are not responsible for the accuracy of news releases posted to EurekAlert! by contributing institutions or for the use of any information through the EurekAlert system.

[Print](#) [E-Mail](#)